



TITLE:

Some Families of Analytic Functions of Complex Order(Study on Calculus Operators in Univalent Function Theory)

AUTHOR(S):

Srivastava, H.M.

CITATION:

Srivastava, H.M.. Some Families of Analytic Functions of Complex Order(Study on Calculus Operators in Univalent Function Theory). 数理解析研究所講究録 2007, 1538: 82-97

ISSUE DATE:

2007-02

URL:

<http://hdl.handle.net/2433/59042>

RIGHT:

Some Families of Analytic Functions of Complex Order

H. M. Srivastava

Department of Mathematics and Statistics

University of Victoria

Victoria, British Columbia V8W 3P4, Canada

E-Mail: harimsri@math.uvic.ca

Abstract

Our *main* objective in this lecture is to present some interesting recent developments concerning inclusion relationships, coefficient bounds, and neighborhood properties associated with certain families of univalent and p -valent analytic functions of *complex* order. Some of the various analytic and multivalent function classes, which are considered in this lecture, are defined by means of the familiar Ruscheweyh derivative operator and its suitably *extended* version applicable to p -valently analytic functions. Several corollaries and consequences of the main results, including relationships with known results, will also be considered briefly.

2000 Mathematics Subject Classification. Primary 30C45; Secondary 05A10, 30A10.

Key Words and Phrases. Analytic functions, univalent and multivalent functions, Hadamard product (or convolution), Ruscheweyh derivative operators, starlike functions, convex functions, (n, δ) -neighborhoods of univalent and p -valent analytic functions, inclusion relationships, coefficient bounds.

H. M. Srivastava

1. Introduction, Definitions and Preliminaries

Let $\mathcal{T}(n, p)$ denote the class of (*normalized*) functions f of the form:

$$f(z) = z^p - \sum_{k=n+p}^{\infty} a_k z^k \quad (1.1)$$

$$(a_k \geq 0; k \in \mathbb{N} \setminus \{1, \dots, n+p-1\}; n, p \in \mathbb{N}; \mathbb{N} := \{1, 2, 3, \dots\}),$$

which are *analytic* and *p-valent* in the *open unit disk*

$$\mathbb{U} := \{z : z \in \mathbb{C} \text{ and } |z| < 1\}.$$

Throughout this presentation, we shall make use of the following *simplified* notations:

$$\mathcal{T}(n, 1) =: \mathcal{T}(n), \mathcal{T}(1, p) =: \mathcal{T}_p \text{ and } \mathcal{T}(1, 1) = \mathcal{T}_1 = \mathcal{T}(1) =: \mathcal{T}.$$

Following the earlier investigations by Goodman [10] and Ruscheweyh [18], we *first* define the (n, δ) -neighborhood of a function $f \in \mathcal{T}(n)$ by (see also [2], [3], [4], and [20])

$$N_{n,\delta}(f) := \left\{ g : g \in \mathcal{T}(n), g(z) = z - \sum_{k=n+1}^{\infty} b_k z^k \text{ and } \sum_{k=n+1}^{\infty} k|a_k - b_k| \leq \delta \right\}. \quad (1.2)$$

In particular, for the *identity* function

$$e(z) = z, \quad (1.3)$$

we immediately have

$$N_{n,\delta}(e) := \left\{ g : g \in \mathcal{T}(n), g(z) = z - \sum_{k=n+1}^{\infty} b_k z^k \text{ and } \sum_{k=n+1}^{\infty} k|b_k| \leq \delta \right\}. \quad (1.4)$$

The above concept of (n, δ) -neighborhoods was extended and applied recently to families of *analytically* multivalent functions by Altıntaş *et al.* [6] and to families of *meromorphically* multivalent functions by Liu and Srivastava ([12] and [13]) (see also the more recent works [17] and [23]). Thus, more generally, we can also define the (n, δ) -neighborhood of a function $f(z) \in \mathcal{T}(n, p)$ ($p \in \mathbb{N}$) by means of the following equation:

$$\mathcal{N}_{n,\delta}(f; p) := \left\{ g : g \in \mathcal{T}(n, p) : g(z) = z^p - \sum_{k=n+p}^{\infty} b_k z^k \text{ and } \sum_{k=n+p}^{\infty} k|a_k - b_k| \leq \delta \right\}, \quad (1.5)$$

so that, obviously,

$$\mathcal{N}_{n,\delta}(h; p) := \left\{ g : g \in \mathcal{T}(n, p) : g(z) = z^p - \sum_{k=n+p}^{\infty} b_k z^k \text{ and } \sum_{k=n+p}^{\infty} k|b_k| \leq \delta \right\}, \quad (1.6)$$

where [cf. Equation (1.3) above]

$$h(z) = z^p \quad (p \in \mathbb{N}) \quad (1.7)$$

Some Families of Analytic Functions of Complex Order

denotes the corresponding *identity* function.

In Sections 2 and 3 of this presentation, we propose to investigate the (n, δ) -neighborhoods of several subclasses of the class $\mathcal{T}(n)$ of *normalized* analytic and univalent functions in \mathbf{U} with negative and missing coefficients, which are introduced here by making use of the Ruscheweyh derivative operator defined by (1.14) or (1.15) below. The rest of this paper deals mainly with the coefficient bounds and inclusion relationships involving the (n, δ) -neighborhoods $N_{n,\delta}(h; p)$ and $N_{n,\delta}(f; p)$ for two other subclasses of the function class $\mathcal{T}(n, p)$, which are introduced in Section 4 of this presentation.

First of all, we say that a function $f \in \mathcal{T}(n)$ is *starlike of complex order* γ ($\gamma \in \mathbb{C} \setminus \{0\}$), that is, $f \in \mathcal{S}_n^*(\gamma)$, if it also satisfies the following inequality:

$$\Re \left(1 + \frac{1}{\gamma} \left[\frac{zf'(z)}{f(z)} - 1 \right] \right) > 0 \quad (z \in \mathbf{U}; \gamma \in \mathbb{C} \setminus \{0\}). \quad (1.8)$$

Furthermore, a function $f \in \mathcal{T}(n)$ is said to be *convex of complex order* γ ($\gamma \in \mathbb{C} \setminus \{0\}$), that is, $f \in \mathcal{C}_n(\gamma)$, if it also satisfies the following inequality:

$$\Re \left(1 + \frac{1}{\gamma} \left[\frac{zf''(z)}{f'(z)} \right] \right) > 0 \quad (z \in \mathbf{U}; \gamma \in \mathbb{C} \setminus \{0\}). \quad (1.9)$$

The classes $\mathcal{S}_n^*(\gamma)$ and $\mathcal{C}_n(\gamma)$ stem essentially from the classes of starlike and convex functions of *complex* order, which were considered earlier by Nasr and Aouf [15] and Wiatrowski [24], respectively (see also [5], [7], [8], [14] and [16] and the relevant other citations in *each* of these works).

Let $\mathcal{S}_n(\gamma, \alpha, \mu, \beta)$ denote the subclass of the function class $\mathcal{T}(n)$ consisting of functions $f(z)$ which satisfy the following inequality:

$$\left| \frac{1}{\gamma} \left(\frac{\alpha\mu z^3 f'''(z) + (2\alpha\mu + \alpha - \mu)z^2 f''(z) + zf'(z)}{\alpha\mu z^2 f''(z) + (\alpha - \mu)zf'(z) + (1 - \alpha + \mu)f(z)} - 1 \right) \right| < \beta \quad (1.10)$$

$$(z \in \mathbf{U}; \gamma \in \mathbb{C} \setminus \{0\}; 0 \leq \mu \leq \alpha; 0 < \beta \leq 1).$$

Suppose also that $\mathcal{R}_n(\gamma, \alpha, \mu, \beta)$ denotes the subclass of the function class $\mathcal{T}(n)$ consisting of functions $f(z)$ which satisfy the the following inequality:

$$\left| \frac{1}{\gamma} (\alpha\mu z^2 f'''(z) + (2\alpha\mu + \alpha - \mu)zf''(z) + f'(z) - 1) \right| < \beta \quad (1.11)$$

$$(z \in \mathbf{U}; \gamma \in \mathbb{C} \setminus \{0\}; 0 \leq \mu \leq \alpha; 0 < \beta \leq 1).$$

The classes $\mathcal{S}_n(\gamma, \alpha, \mu, \beta)$ and $\mathcal{R}_n(\gamma, \alpha, \mu, \beta)$ were studied recently by Orhan and Kamali [16].

Next, for the functions $f_j(z)$ ($j = 1, 2$) given by

$$f_j(z) = z + \sum_{k=2}^{\infty} a_{k,j} z^k \quad (j = 1, 2), \quad (1.12)$$

H. M. Srivastava

we denote by $(f_1 \star f_2)(z)$ the Hadamard product (or convolution) of $f_1(z)$ and $f_2(z)$, defined by

$$(f_1 \star f_2)(z) := z + \sum_{k=2}^{\infty} a_{k,1} a_{k,2} z^k =: (f_2 \star f_1)(z). \quad (1.13)$$

Thus the Ruscheweyh derivative operator

$$D^\lambda : \mathcal{T} \rightarrow \mathcal{T} \quad (\mathcal{T} := \mathcal{T}(1) = \mathcal{T}_1 = \mathcal{T}(1, 1))$$

is defined by

$$D^\lambda f(z) := \frac{z}{(1-z)^{\lambda+1}} \star f(z) \quad (\lambda > -1; f \in \mathcal{T}). \quad (1.14)$$

or, equivalently, by

$$D^\lambda f(z) := z - \sum_{k=2}^{\infty} \binom{\lambda+k-1}{k-1} a_k z^k \quad (\lambda > -1; f \in \mathcal{T}) \quad (1.15)$$

for a function $f \in \mathcal{T}$ of the form (1.1). Here, and in what follows, we make use of the following standard notation for a binomial coefficient:

$$\binom{\kappa}{n} := \frac{\kappa(\kappa-1)\cdots(\kappa-n+1)}{n!} \quad (\kappa \in \mathbb{C}; n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}). \quad (1.16)$$

In particular, we have

$$D^n f(z) = \frac{z(z^{n-1}f(z))^{(n)}}{n!} \quad (n \in \mathbb{N}_0). \quad (1.17)$$

Finally, in terms of the Ruscheweyh derivative operator D^λ ($\lambda > -1$) defined by (1.14) or (1.15) above, let $\mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta)$ denote the subclass of the function class $\mathcal{T}(n)$ consisting of functions $f(z)$ which satisfy the following inequality:

$$\left| \frac{1}{\gamma} \left(\frac{\alpha\mu z^3 (D^\lambda f(z))''' + (2\alpha\mu + \alpha - \mu)z^2 (D^\lambda f(z))'' + z(D^\lambda f(z))'}{\alpha\mu z^2 (D^\lambda f(z))'' + (\alpha - \mu)z(D^\lambda f(z))' + (1 - \alpha + \mu)D^\lambda f(z)} - 1 \right) \right| < \beta \quad (1.18)$$

$$(z \in \mathbb{U}; \gamma \in \mathbb{C} \setminus \{0\}; \lambda > -1; 0 < \beta \leq 1; 0 \leq \mu \leq \alpha).$$

Also let $\mathcal{R}_n(\gamma, \lambda, \alpha, \mu, \beta)$ denote the subclass of the function class $\mathcal{T}(n)$ consisting of functions $f(z)$ which satisfy the following inequality:

$$\left| \frac{1}{\gamma} \left(\alpha\mu z^2 (D^\lambda f(z))''' + (2\alpha\mu + \alpha - \mu)z(D^\lambda f(z))'' + (D^\lambda f(z))' - 1 \right) \right| < \beta \quad (1.19)$$

$$(z \in \mathbb{U}; \gamma \in \mathbb{C} \setminus \{0\}; \lambda > -1; 0 < \beta \leq 1; 0 \leq \mu \leq \alpha).$$

Various *further* subclasses of the function class $\mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta)$ with

$$\gamma = 1 \quad \text{and} \quad \alpha = \mu = 0 \quad (1.20)$$

Some Families of Analytic Functions of Complex Order

were studied in many earlier works (*cf.*, *e.g.*, [9], [11], [21] and [22]; see also the references cited in each of these earlier works). Clearly, in these cases of (for example) the class $\mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta)$, we have the following relationships:

$$\mathcal{S}_n(\gamma, 0, 0, 0, 1) \subset \mathcal{S}_n^*(\gamma) \quad \text{and} \quad \mathcal{S}_n(\gamma, 0, 1, 0, 1) \subset \mathcal{C}_n(\gamma) \quad (1.21)$$

$$(n \in \mathbb{N}; \gamma \in \mathbb{C} \setminus \{0\}).$$

2. Inclusion Relationships Involving the (n, δ) -Neighborhood $N_{n,\delta}(e)$

In our investigation of the inclusion relationships involving the (n, δ) -neighborhood $N_{n,\delta}(e)$ defined by (1.4), we shall require the following lemmas.

Lemma 1. *Let $f \in \mathcal{T}(n)$ be defined by (1.1) (with $p = 1$). Then f is in the class $\mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta)$ if and only if*

$$\sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} \eta(k) a_k \leq \beta|\gamma|, \quad (2.1)$$

where

$$\eta = \eta(k) := \left(\alpha \mu k^3 + (\alpha - \mu - 2\alpha\mu + \alpha\mu\beta|\gamma|)k^2 \right. \\ \left. + (\alpha\mu - 2\alpha - 2\mu + 1 + (\alpha - \mu - \alpha\mu)\beta|\gamma|)k + (1 - \alpha + \mu)(\beta|\gamma| - 1) \right).$$

Proof. We first suppose that $f \in \mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta)$. Then, by appealing to the condition (1.18), we readily find that

$$\Re \left(\frac{\alpha \mu z^3 (D^\lambda f(z))''' + (2\alpha\mu + \alpha - \mu)z^2 (D^\lambda f(z))'' + z(D^\lambda f(z))'}{\alpha \mu z^2 (D^\lambda f(z))'' + (\alpha - \mu)z(D^\lambda f(z))' + (1 - \alpha + \mu)D^\lambda f(z)} - 1 \right) \\ > -\beta|\gamma| \quad (z \in \mathbb{U}) \quad (2.2)$$

or, equivalently, that

$$\Re \left(\frac{- \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} [\alpha \mu k^3 + (\alpha - \mu - 2\alpha\mu)k^2 + (\alpha\mu - 2\alpha + 2\mu + 1)k - (1 - \alpha + \mu)] a_k z^k}{z - \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} [\alpha \mu k^2 + (\alpha - \mu - \alpha\mu)k + (1 - \alpha + \mu)] a_k z^k} \right) \\ > -\beta|\gamma| \quad (z \in \mathbb{U}), \quad (2.3)$$

where we have made use of the explicit representation (1.15) and the definition (1.1) (with $p = 1$). We now choose values of z on the real axis and let $z \rightarrow 1-$ through *real* values. Then the inequality (2.3) immediately yields the desired condition (2.1).

H. M. Srivastava

Conversely, by applying the hypothesis (2.1) and letting $|z| = 1$, we find that

$$\begin{aligned}
 & \left| \frac{\alpha\mu z^3 (D^\lambda f(z))''' + (2\alpha\mu + \alpha - \mu)z^2 (D^\lambda f(z))''' + z(D^\lambda f(z))'}{\alpha\mu z^2 (D^\lambda f(z))'' + (\alpha - \mu)z(D^\lambda f(z))' + (1 - \alpha + \mu)D^\lambda f(z)} - 1 \right| \\
 &= \left| \frac{\sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} [\alpha\mu k^3 + (\alpha - \mu - 2\alpha\mu)k^2 + (\alpha\mu - 2\alpha + 2\mu + 1)k - (1 - \alpha + \mu)] a_k z^k}{1 - \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} [\alpha\mu k^2 + (\alpha - \mu - \alpha\mu)k + (1 - \alpha + \mu)] a_k z^k} \right| \\
 &\leq \frac{\beta|r| \left[1 - \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} [\alpha\mu k^2 + (\alpha - \mu - \alpha\mu)k + (1 - \alpha + \mu)] a_k \right]}{1 - \sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} [\alpha\mu k^2 + (\alpha - \mu - \alpha\mu)k + (1 - \alpha + \mu)] a_k} \\
 &\leq \beta|\gamma|. \tag{2.4}
 \end{aligned}$$

Hence, by the *maximum modulus principle*, we have

$$f \in \mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta),$$

which evidently completes the proof of Lemma 1.

Similarly, we can prove the following result.

Lemma 2. *Let the function $f \in \mathcal{T}(n)$ be defined by (1.1) (with $p = 1$). Then f is in the class $\mathcal{R}_n(\gamma, \lambda, \alpha, \mu, \beta)$ if only if*

$$\sum_{k=n+1}^{\infty} \binom{\lambda+k-1}{k-1} [\alpha\mu k^3 + (\alpha - \mu - \alpha\mu)k^2 + (1 - \alpha + \mu)k] a_k \leq \beta|\gamma|. \tag{2.5}$$

Remark 1. A special case of Lemma 1 when

$$n = 1, \mu = \alpha = 0, \gamma = 1 \quad \text{and} \quad \beta = 1 - c \quad (0 \leq c < 1)$$

was given by Ahuja [1]. Furthermore, in Lemma 1 with

$$n = 1, \mu = \alpha = 0, \gamma = 1 \quad \text{and} \quad \beta = 1 - c \quad (0 \leq c < 1),$$

if we set

$$\lambda = 0 \quad \text{and} \quad \lambda = 1.$$

we obtain the relatively more familiar results of Silverman [19].

Our first main result is given by Theorem 1 below.

Theorem 1. *If*

$$\delta := \frac{(n+1)\beta|\gamma|}{\binom{\lambda+n}{n}^\rho}, \tag{2.6}$$

Some Families of Analytic Functions of Complex Order

then

$$\mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta) \subset N_{n,\delta}(e), \quad (2.7)$$

where

$$\begin{aligned} \rho := & \left[\alpha\mu(n+1)^3 + (\alpha\mu\beta|\gamma| + \alpha - \mu - 2\alpha\mu)(n+1)^2 \right. \\ & + ((\alpha - \mu - \alpha\mu)\beta|\gamma| + 1 - 2\alpha + 2\mu + \alpha\mu)(n+1) \\ & \left. + (1 - \alpha + \mu)(\beta|\gamma| - 1) \right]. \end{aligned} \quad (2.8)$$

Proof. For a function $f \in \mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta)$ of the form (1.1) (with $p = 1$) and for ρ defined already by (2.8), Lemma 1 immediately yields

$$\binom{\lambda+n}{n} \rho \sum_{k=n+1}^{\infty} a_k \leq \beta|\gamma|,$$

so that

$$\sum_{k=n+1}^{\infty} a_k \leq \frac{\beta|\gamma|}{\binom{\lambda+n}{n} \rho}. \quad (2.9)$$

On the other hand, we also find from (2.1) that

$$\binom{\lambda+n}{n} \tau \sum_{k=n+1}^{\infty} k a_k \leq \beta|\gamma|,$$

where

$$\begin{aligned} \tau = & \left[\alpha\mu(n+1)^2 + (\alpha\mu\beta|\gamma| + \alpha - \mu - 2\alpha\mu)(n+1) \right. \\ & + ((\alpha - \mu - \alpha\mu)\beta|\gamma| + 1 - 2\alpha + 2\mu + \alpha\mu) \\ & \left. + \left(\frac{(1 - \alpha + \mu)(\beta|\gamma| - 1)}{n+1} \right) \right], \end{aligned} \quad (2.10)$$

that is, that

$$\sum_{k=n+1}^{\infty} k a_k \leq \frac{\beta|\gamma|(n+1)}{\binom{\lambda+n}{n} \rho} := \delta, \quad (2.11)$$

which, in view of the definition (1.4), proves Theorem 1.

In a similar manner, by applying Lemma 2 instead of Lemma 1, we can prove Theorem 2 below.

H. M. Srivastava

Theorem 2. *If*

$$\delta := \frac{\beta|\gamma|}{\binom{\lambda+n}{n} [\alpha\mu(n+1)^2 + (\alpha - \mu - \alpha\mu)(n+1) + (1 - \alpha + \mu)]}, \quad (2.12)$$

then

$$\mathcal{R}_n(\gamma, \lambda, \alpha, \mu, \beta) \subset N_{n,\delta}(e).$$

3. Neighborhood Properties for the Function Classes

$$\mathcal{S}_n^{(b)}(\gamma, \lambda, \alpha, \mu, \beta) \quad \text{and} \quad \mathcal{R}_n^{(b)}(\gamma, \lambda, \alpha, \mu, \beta)$$

In this section, we determine the neighborhood for each of the function classes

$$\mathcal{S}_n^{(b)}(\gamma, \lambda, \alpha, \mu, \beta) \quad \text{and} \quad \mathcal{R}_n^{(b)}(\gamma, \lambda, \alpha, \mu, \beta),$$

which we define here as follows.

Definition 1. A function $f \in \mathcal{T}(n)$ is said to be in the class $\mathcal{S}_n^{(b)}(\gamma, \lambda, \alpha, \mu, \beta)$ if there exists a function

$$g \in \mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta)$$

such that the following inequality holds true:

$$\left| \frac{f(z)}{g(z)} - 1 \right| < 1 - b \quad (z \in \mathbf{U}; 0 \leq b < 1). \quad (3.1)$$

Definition 2. A function $f \in \mathcal{T}(n)$ is said to be in the class $\mathcal{R}_n^{(b)}(\gamma, \lambda, \alpha, \mu, \beta)$ if there exists a function

$$g \in \mathcal{R}_n(\gamma, \lambda, \alpha, \mu, \beta)$$

such that the inequality (3.1) holds true.

Theorem 3. *If* $g \in \mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta)$ *and*

$$b = 1 - \frac{\binom{\lambda+n}{n} \delta \rho}{(n+1) \left[\binom{\lambda+n}{n} \rho - \beta|\gamma| \right]}, \quad (3.2)$$

then

$$N_{n,\delta}(g) \subset \mathcal{S}_n^{(b)}(\gamma, \lambda, \alpha, \mu, \beta), \quad (3.3)$$

where ρ is given already by (2.8).

Proof. Assuming that $f \in N_{n,\delta}(g)$, we find from the definition (1.2) that

$$\sum_{k=n+1}^{\infty} k|a_k - b_k| \leq \delta, \quad (3.4)$$

Some Families of Analytic Functions of Complex Order

which readily implies the following coefficient inequality:

$$\sum_{k=n+1}^{\infty} |a_k - b_k| \leq \frac{\delta}{n+1} \quad (n \in \mathbb{N}). \quad (3.5)$$

Since $g \in \mathcal{S}_n(\gamma, \lambda, \alpha, \mu, \beta)$, we have [cf. Equation (2.9)].

$$\sum_{k=n+1}^{\infty} b_k = \frac{\beta|\gamma|}{\binom{\lambda+n}{n} \rho}, \quad (3.6)$$

so that

$$\begin{aligned} \left| \frac{f(z)}{g(z)} - 1 \right| &< \frac{\sum_{k=n+1}^{\infty} |a_k - b_k|}{1 - \sum_{k=n+1}^{\infty} b_k} \\ &\leq \frac{\delta}{n+1} \cdot \frac{\binom{\lambda+n}{n} \delta \rho}{\left[\binom{\lambda+n}{n} \rho - \beta|\gamma| \right]} =: 1 - b, \end{aligned} \quad (3.7)$$

provided that b is given precisely by (3.2). Thus, by Definition 1, we conclude that

$$f \in \mathcal{S}_n^{(b)}(\gamma, \lambda, \alpha, \mu, \beta)$$

for b given by (3.2). This evidently completes the proof of Theorem 3.

The proof of Theorem 4 below is much akin to that of Theorem 3, and so the details involved are being omitted here.

Theorem 4. If $g \in \mathcal{R}_n(\gamma, \lambda, \alpha, \mu, \beta)$ and

$$b = 1 - \frac{\binom{\lambda+n}{n} \delta [\alpha\mu(n+1)^3 + (\alpha - \mu - \alpha\mu)(n+1)^2 + (1 - \alpha + \mu)(n+1)]}{(n+1) \left[\binom{\lambda+n}{n} [\alpha\mu(n+1)^3 + (\alpha - \mu - \alpha\mu)(n+1)^2 + (1 - \alpha + \mu)(n+1)] - \beta|\gamma| \right]}, \quad (3.8)$$

then

$$N_{n,\delta}(g) \subset \mathcal{R}_n^{(b)}(\gamma, \lambda, \alpha, \mu, \beta). \quad (3.9)$$

Remark 2. A special case of Theorem 3 when $\alpha = \mu = 0$ was proven recently by Murugusundaramoorthy and Srivastava [14].

H. M. Srivastava

4. A Set of Coefficient Bounds for the Function Classes

$$\mathcal{H}_{n,m}^p(\lambda, b) \quad \text{and} \quad \mathcal{L}_{n,m}^p(\lambda, b; \mu)$$

With a view to introducing the function classes

$$\mathcal{H}_{n,m}^p(\lambda, b) \quad \text{and} \quad \mathcal{L}_{n,m}^p(\lambda, b; \mu),$$

we begin by considering the Hadamard product (or convolution) of the function $f \in \mathcal{T}(n, p)$ given by (1.1) and the function $g \in \mathcal{T}(n, p)$ given by

$$g(z) = z^p - \sum_{k=n+p}^{\infty} b_k z^k \quad (b_k \geq 0; n, p \in \mathbb{N}), \quad (4.1)$$

which is defined (*as usual*) by

$$(f * g)(z) := z^p + \sum_{k=n+p}^{\infty} a_k b_k z^k =: (g * f)(z). \quad (4.2)$$

We next introduce an *extended* linear derivative operator of the Ruscheweyh type given already by (1.11) or (1.12) above:

$$\mathcal{D}^{\lambda, p} : \mathcal{T}_p \rightarrow \mathcal{T}_p \quad (\mathcal{T}_p := \mathcal{T}(1, p)),$$

which is defined here by the following convolution:

$$\mathcal{D}^{\lambda, p} f(z) := \frac{z^p}{(1-z)^{\lambda+p}} * f(z) \quad (\lambda > -p; f \in \mathcal{T}_p). \quad (4.3)$$

In terms of the binomial coefficients in (1.16), we can rewrite (4.3) as follows:

$$\mathcal{D}^{\lambda, p} f(z) = z^p - \sum_{k=1+p}^{\infty} \binom{\lambda+k-1}{k-p} a_k z^k \quad (\lambda > -p; f \in \mathcal{T}_p). \quad (4.4)$$

In particular, when $\lambda = n$ ($n \in \mathbb{N}$), it is easily observed from (4.3) and (4.4) that

$$\mathcal{D}^{n, p} f(z) = \frac{z^p (z^{n-p} f(z))^{(n)}}{n!} \quad (n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}; p \in \mathbb{N}), \quad (4.5)$$

so that

$$\mathcal{D}^{1, p} f(z) = (1-p)f(z) + z f'(z), \quad (4.6)$$

$$\mathcal{D}^{2, p} f(z) = \frac{(1-p)(2-p)}{2!} f(z) + (2-p)z f'(z) + \frac{z^2}{2!} f''(z), \quad (4.7)$$

and so on. In fact, by comparing the definitions (1.14) and (4.3), we readily have

$$\mathcal{D}^{\lambda, 1} f(z) =: D^{\lambda} f(z) \quad (\lambda > -1; f \in \mathcal{T}). \quad (4.8)$$

By using this *extended* Ruscheweyh derivative operator

$$\mathcal{D}^{\lambda, p} f(z) \quad (\lambda > -p; p \in \mathbb{N})$$

Some Families of Analytic Functions of Complex Order

given by (4.4), we now introduce a new subclass $\mathcal{H}_{n,m}^p(\lambda, b)$ of the p -valently analytic function class $\mathcal{T}(n, p)$, which includes functions $f(z)$ satisfying the following inequality:

$$\left| \frac{1}{b} \left(\frac{z(\mathcal{D}^{\lambda,p} f(z))^{(m+1)}}{(\mathcal{D}^{\lambda,p} f(z))^{(m)}} - (p-m) \right) \right| < 1 \quad (4.9)$$

($z \in U$; $p \in \mathbb{N}$; $m \in \mathbb{N}_0$; $\lambda \in \mathbb{R}$; $p > \max(m, -\lambda)$; $b \in \mathbb{C} \setminus \{0\}$).

We also denote by $\mathcal{L}_{n,m}^p(\lambda, b; \mu)$ the subclass of $\mathcal{T}(n, p)$ consisting of functions $f(z)$ which satisfy the inequality (4.10) below:

$$\left| \frac{1}{b} \left(p(1-\mu) \left(\frac{\mathcal{D}^{\lambda,p} f(z)}{z} \right)^{(m)} + \mu (\mathcal{D}^{\lambda,p} f(z))^{(m+1)} - (p-m) \right) \right| < p-m \quad (4.10)$$

($z \in U$; $p \in \mathbb{N}$; $m \in \mathbb{N}_0$; $\lambda \in \mathbb{R}$; $p > \max(m, -\lambda)$; $\mu \geq 0$; $b \in \mathbb{C} \setminus \{0\}$).

Our definitions of the function classes

$$\mathcal{H}_{n,m}^p(\lambda, b) \quad \text{and} \quad \mathcal{L}_{n,m}^p(\lambda, b; \mu)$$

are motivated essentially by two earlier investigations [4] and [14], in each of which further details and references to other closely-related subclasses can be found. In particular, in our definition of the function class $\mathcal{L}_{n,m}^p(\lambda, b; \mu)$ involving the inequality (1.13), we have relaxed the parametric constraint $0 \leq \mu \leq 1$, which was imposed earlier by Murugusundaramoorthy and Srivastava [14, p. 3, Equation (1.14)] (see also Remark 5 below).

We now prove the following results which yield the coefficient inequalities for functions in the subclasses (see also [17])

$$\mathcal{H}_{n,m}^p(\lambda, b) \quad \text{and} \quad \mathcal{L}_{n,m}^p(\lambda, b; \mu).$$

Theorem 5. *Let $f(z) \in \mathcal{T}(n, p)$ be given by (1.1). Then $f(z) \in \mathcal{H}_{n,m}^p(\lambda, b)$ if and only if*

$$\sum_{k=n+p}^{\infty} \binom{\lambda+k-1}{k-p} \binom{k}{m} (k+|b|-p) a_k \leq |b| \binom{p}{m}. \quad (4.11)$$

Proof. Let a function $f(z)$ of the form (1.1) belong to the class $\mathcal{H}_{n,m}^p(\lambda, b)$. Then, in view of (4.4), (4.9) yields the following inequality:

$$\Re \left(\frac{\sum_{k=n+p}^{\infty} \binom{\lambda+k-1}{k-p} \binom{k}{m} (p-k) z^{k-p}}{\binom{p}{m} - \sum_{k=n+p}^{\infty} \binom{\lambda+k-1}{k-p} \binom{k}{m} z^{k-p}} \right) > -|b| \quad (z \in U). \quad (4.12)$$

Putting $z = r$ ($0 \leq r < 1$) in (4.12), we observe that the expression in the denominator on the left-hand side of (2.2) is positive for $r = 0$ and also for all r ($0 < r < 1$). Thus, by letting $r \rightarrow 1-$ through real values, (4.12) leads us to the desired assertion (2.1) of Theorem 5.

H. M. Srivastava

Conversely, by applying (4.11) and setting $|z| = 1$, we find by using (4.4) that

$$\begin{aligned}
 & \left| \frac{z (\mathcal{D}^{\lambda,p} f(z))^{(m+1)}}{(\mathcal{D}^{\lambda,p} f(z))^{(m)}} - (p-m) \right| \\
 &= \left| \frac{\sum_{k=n+p}^{\infty} \binom{\lambda+k-1}{k-p} \binom{k}{m} (p-k) z^{k-m}}{\binom{p}{m} z^{p-m} - \sum_{k=n+p}^{\infty} \binom{\lambda+k-1}{k-p} \binom{k}{m} z^{k-m}} \right| \\
 &\leq \frac{|b| \left[\binom{p}{m} - \sum_{k=n+p}^{\infty} \binom{\lambda+k-1}{k-p} \binom{k}{m} a_k \right]}{\binom{p}{m} - \sum_{k=n+p}^{\infty} \binom{\lambda+k-1}{k-p} \binom{k}{m} a_k} = |b|. \quad (4.13)
 \end{aligned}$$

Hence, by the *maximum modulus principle* once again, we infer that $f(z) \in \mathcal{H}_{n,m}^p(\lambda, b)$, which completes the proof of Theorem 5.

Remark 3. In the special case when

$$m = 0, \quad p = 1, \quad \text{and} \quad b = \beta\gamma \quad (0 < \beta \leq 1; \gamma \in \mathbb{C} \setminus \{0\}), \quad (4.14)$$

Theorem 1 corresponds to a result given earlier by Murugusundaramoorthy and Srivastava [14, p. 3, Lemma 1].

By using the same arguments as in the proof of Theorem 5, we can establish Theorem 6 below.

Theorem 6. Let $f(z) \in \mathcal{T}(n, p)$ be given by (1.1). Then $f(z) \in \mathcal{L}_{n,m}^p(\lambda, b; \mu)$ if and only if

$$\begin{aligned}
 & \sum_{k=n+p}^{\infty} \binom{\lambda+k-1}{k-p} \binom{k-1}{m} [\mu(k-1) + 1] a_k \\
 & \leq (p-m) \left[\frac{|b|-1}{m!} + \binom{p}{m} \right]. \quad (4.15)
 \end{aligned}$$

Remark 4. Making use of the same parametric substitutions as mentioned above in (2.3), Theorem 2 yields another known result due to Murugusundaramoorthy and Srivastava [14, p. 4, Lemma 2].

Some Families of Analytic Functions of Complex Order

5. Inclusion Relationships Involving the (n, δ) -Neighborhood $\mathcal{N}_{n,\delta}(h; p)$

In this section, we establish several inclusion relationships for the function classes

$$\mathcal{H}_{n,m}^p(\lambda, b) \quad \text{and} \quad \mathcal{L}_{n,m}^p(\lambda, b; \mu)$$

involving the (n, δ) -neighborhood defined by (1.6).

Theorem 7. *If*

$$\delta = \frac{(n+p)|b| \binom{p}{m}}{(n+|b|) \binom{\lambda+n+p-1}{n} \binom{n+p}{m}} \quad (p > |b|), \quad (5.1)$$

then

$$\mathcal{H}_{n,m}^p(\lambda, b) \subset \mathcal{N}_{n,\delta}(h; p). \quad (5.2)$$

Proof. Let $f(z) \in \mathcal{H}_{n,m}^p(\lambda, b)$. Then, in view of the assertion (4.11) of Theorem 5, we have

$$(n+|b|) \binom{\lambda+n+p-1}{n} \binom{n+p}{m} \sum_{k=n+p}^{\infty} a_k \leq |b| \binom{p}{m}. \quad (5.3)$$

This yields

$$\sum_{k=n+p}^{\infty} a_k \leq \frac{|b| \binom{p}{m}}{(n+|b|) \binom{\lambda+n+p-1}{n} \binom{n+p}{m}}. \quad (5.4)$$

Applying the assertion (4.11) of Theorem 5 again, in conjunction with (5.4), we observe that

$$\begin{aligned} & \binom{\lambda+n+p-1}{n} \binom{n+p}{m} \sum_{k=n+p}^{\infty} k a_k \\ & \leq |b| \binom{p}{m} + (p-|b|) \binom{\lambda+n+p-1}{n} \binom{n+p}{m} \sum_{k=n+p}^{\infty} a_k \\ & \leq |b| \binom{p}{m} + (p-|b|) \binom{\lambda+n+p-1}{n} \binom{n+p}{m} \\ & \quad \cdot \frac{|b| \binom{p}{m}}{(n+|b|) \binom{\lambda+n+p-1}{n} \binom{n+p}{m}} \\ & = |b| \binom{p}{m} \left(\frac{n+p}{n+|b|} \right). \end{aligned}$$

H. M. Srivastava

Hence we have

$$\sum_{k=n+p}^{\infty} ka_k \leq \frac{|b|(n+p) \binom{p}{m}}{(n+|b|) \binom{\lambda+n+p-1}{n} \binom{n+p}{m}} =: \delta \quad (p > |b|), \quad (5.5)$$

which, by virtue of (1.6), establishes the inclusion relation (5.2) of Theorem 7.

In an analogous manner, by applying the assertion (4.15) of Theorem 6 instead of the assertion (4.11) of Theorem 5 to functions in the class $\mathcal{L}_{n,m}^p(\lambda, b; \mu)$, we can prove the following inclusion relationship.

Theorem 8. *If*

$$\delta = \frac{(p-m)(n+p) \left[\frac{|b|-1}{m!} + \binom{p}{m} \right]}{[\mu(n+p-1)+1] \binom{\lambda+n+p-1}{n} \binom{n+p}{m}} \quad (\mu > 1), \quad (5.6)$$

then

$$\mathcal{L}_{n,m}^p(\lambda, b; \mu) \subset \mathcal{N}_{n,\delta}(h; p).$$

Remark 5. Applying the parametric substitutions listed in (4.14), Theorems 7 and 8 would yield the known results due to Murugusundaramoorthy and Srivastava [14, p. 4, Theorem 1; p. 5, Theorem 2]. Incidentally, just as we indicated in Section 4 above, the condition $\mu > 1$ is needed in the proof of one of these known results [14, p. 5, Theorem 2]. This implies that the constraint $0 \leq \mu \leq 1$ in [14, p. 3, Equation (1.14)] should be replaced by the less stringent constraint $\mu \geq 0$ (see also [17, p. 5, Remark 3]).

6. Further Neighborhood Properties Involving $\mathcal{N}_{n,\delta}(f; p)$

In this last section, we determine the neighborhood properties for each of the following (*slightly modified*) function classes:

$$\mathcal{H}_{n,m}^{p,\alpha}(\lambda, b) \quad \text{and} \quad \mathcal{L}_{n,m}^{p,\alpha}(\lambda, b; \mu).$$

Here, by definition, the class $\mathcal{H}_{n,m}^{p,\alpha}(\lambda, b)$ consists of functions $f(z) \in \mathcal{T}(n, p)$ for which there exists another function

$$g(z) \in \mathcal{H}_{n,m}^p(\lambda, b)$$

such that

$$\left| \frac{f(z)}{g(z)} - 1 \right| < p - \alpha \quad (z \in \mathbb{U}; 0 \leq \alpha < p). \quad (6.1)$$

Analogously, the class $\mathcal{L}_{n,m}^{p,\alpha}(\lambda, b; \mu)$ consists of functions $f(z) \in \mathcal{T}(n, p)$ for which there exists another function

$$g(z) \in \mathcal{L}_{n,m}^p(\lambda, b; \mu)$$

Some Families of Analytic Functions of Complex Order

satisfying the inequality (6.1).

The proofs of the following results (Theorems 9 and 10) involving the neighborhood properties for the classes

$$\mathcal{H}_{n,m}^{p,\alpha}(\lambda, b) \quad \text{and} \quad \mathcal{L}_{n,m}^{p,\alpha}(\lambda, b; \mu)$$

are similar to those given already by Altıntaş *et al.* [4] and, more recently, by Murugusundaramoorthy and Srivastava [14]. We, therefore, choose to skip their proofs here.

Theorem 9. Suppose that

$$g(z) \in \mathcal{H}_{n,m}^p(\lambda, b).$$

Also let

$$\alpha = p - \frac{\delta(n+|b|) \binom{\lambda+n+p-1}{n} \binom{n+p}{m}}{(n+p) \left[(n+|b|) \binom{\lambda+n+p-1}{n+p} \binom{n+p}{m} - |b| \binom{p}{m} \right]}. \quad (6.2)$$

Then

$$\mathcal{N}_{n,\delta}(g; p) \subset \mathcal{H}_{n,m}^{p,\alpha}(\lambda, b).$$

Theorem 10. Suppose that

$$g(z) \in \mathcal{L}_{n,m}^p(\lambda, b; \mu).$$

Also let

$$\alpha = p - \frac{\delta[\mu(n+p-1)+1] \binom{\lambda+n+p-1}{n} \binom{n+p-1}{m}}{(n+p) \left[[\mu(n+p-1)+1] \binom{\lambda+n+p-1}{n} \binom{n+p-1}{m} - (p-m) \left\{ \frac{|b|-1}{m!} + \binom{p}{m} \right\} \right]}. \quad (6.3)$$

Then

$$\mathcal{N}_{n,\delta}(g; p) \subset \mathcal{L}_{n,m}^{p,\alpha}(\lambda, b; \mu).$$

Acknowledgements

It is a great pleasure for me to express my sincere thanks to the members of the Organizing Committee of this *RIMS (Kyoto University) International Short Joint Research Workshop on Calculus Operators in Univalent Function Theory* (especially to Professor Shigeyoshi Owa) for their kind invitation and excellent hospitality. Indeed I am immensely grateful also to many other friends and colleagues in Japan for their having made my visit to Japan in May 2006 a rather pleasant, memorable, and professionally fruitful visit. The present investigation was supported, in part, by the *Natural Sciences and Engineering Research Council of Canada* under Grant OGP0007353.

H. M. Srivastava

References

- [1] O. P. Ahuja, Hadamard products of analytic functions defined by Ruscheweyh derivatives, in *Current Topics in Analytic Function Theory* (H. M. Srivastava and S. Owa, Editors), pp. 13-28, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, 1992.
- [2] O. P. Ahuja and M. Nunokawa, Neighborhoods of analytic functions defined by Ruscheweyh derivatives, *Math. Japon.* **51** (2003), 487-492.
- [3] O. Altıntaş and S. Owa, Neighborhoods of certain analytic functions with negative coefficients, *Internat. J. Math. and Math. Sci.* **19** (1996), 797-800.
- [4] O. Altıntaş, Ö. Özkan and H. M. Srivastava, Neighborhoods of a class of analytic functions with negative coefficients, *Appl. Math. Lett.* **13** (3) (2000), 63-67.
- [5] O. Altıntaş, Ö. Özkan and H. M. Srivastava, Majorization by starlike functions of complex order, *Complex Variables Theory Appl.* **46** (2001), 207-218.
- [6] O. Altıntaş, Ö. Özkan and H. M. Srivastava, Neighborhoods of a certain family of multivalent functions with negative coefficients, *Comput. Math. Appl.* **47** (2004), 1667-1672.
- [7] O. Altıntaş and H. M. Srivastava, Some majorization problems associated with p -valently starlike and convex functions of complex order, *East Asian Math. J.* **17** (2001), 175-183.
- [8] M. K. Aouf, H. M. Hossen and H. E. El-Attar, Certain classes of analytic functions of complex order and type beta with fixed second coefficient, *Math. Sci. Res. Hot-Line* **4** (4) (2000), 31-45.
- [9] P. L. Duren, *Univalent Functions*, A Series of Comprehensive Studies in Mathematics, Vol. **259**, Springer-Verlag, New York, Berlin, Heidelberg and Tokyo, 1983.
- [10] A. W. Goodman, Univalent functions and nonanalytic curves, *Proc. Amer. Math. Soc.* **8** (1957), 598-601.
- [11] A. W. Goodman, *Univalent Functions*, Vol. I, Mariner Publishing Company, Tampa, Florida, 1983.
- [12] J.-L. Liu and H. M. Srivastava, Classes of meromorphically multivalent functions associated with the generalized hypergeometric function, *Math. Comput. Modelling* **39** (2004), 21-34.
- [13] J.-L. Liu and H. M. Srivastava, Subclasses of meromorphically multivalent functions associated with a certain linear operator, *Math. Comput. Modelling* **39** (2004), 35-44.
- [14] G. Murugusundaramoorthy and H. M. Srivastava, Neighborhoods of certain classes of analytic functions of complex order, *J. Inequal. Pure Appl. Math.* **5** (2) (2004), Article 24, 1-8 (electronic).
- [15] M. A. Nasr and M. K. Aouf, Starlike function of complex order, *J. Natur. Sci. Math.* **25** (1985), 1-12.
- [16] H. Orhan and M. Kamali, Starlike, convex and close-to-convex functions of complex order, *Appl. Math. Comput.* **135** (2003), 251-262.
- [17] R. K. Raina and H. M. Srivastava, Inclusion and neighborhood properties of some analytic and multivalent functions, *J. Inequal. Pure Appl. Math.* **7** (1) (2006), Article 5, 1-6 (electronic).
- [18] S. Ruscheweyh, Neighborhoods of univalent functions, *Proc. Amer. Math. Soc.* **81** (1981), 521-527.
- [19] H. Silverman, Univalent functions with negative coefficients, *Proc. Amer. Math. Soc.* **51** (1975), 109-116.
- [20] H. Silverman, Neighborhoods of classes of analytic functions, *Far East J. Math. Sci.* **3** (1995), 165-169.
- [21] H. M. Srivastava and S. Owa (Editors), *Univalent Functions, Fractional Calculus, and Their Applications*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1989.
- [22] H. M. Srivastava and S. Owa (Editors), *Current Topics in Analytic Function Theory*, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, 1992.
- [23] H. M. Srivastava and J. Patel, Some subclasses of multivalent functions involving a certain linear operator, *J. Math. Anal. Appl.* **310** (2005), 209-228.
- [24] P. Wiatrowski, On the coefficients of some family of holomorphic functions, *Zeszyty Nauk. Univ. Łódź Nauk. Mat.-Przyrod.* (2) **39** (1970), 75-85.